

Saturnalia

Colin Wilkin

University College London, London, WC1E 6BT, UK

Abstract

A personal choice is made of the highlights of the physics programme carried out at the Saturne-2 facility over the last twenty years.

It is the end, the end of the meeting and the end of Saturne, and the kind organisers have asked me to say a few words about both events. We have been gathered here in Paris for the last two days to celebrate the work of a whole collection of people and apparatus grouped around an accelerator Saturne-2, which closed down in December. I will not give a political speech but there is an important point which has to be made. The NSF facility at Daresbury was shut in the 1980's, at the height of its productivity, by a committee of physicists who were faced with a budgetary crisis caused by the need to support particle physics at a time of a rising Swiss Franc. When nuclear physicists queried the action, the administration replied that it was a decision taken on physics grounds, which is not the same as a decision taken by physicists under pressure! As a result of this deliberate confusion, which was fed to the media, not only had the nuclear physicists lost their machine, but they felt devalued and insulted into the bargain. Do not let anyone tell you that Saturne could not have continued because there were not interesting problems still to be studied or that Saturne was not competitive for their study — both would be false.

The flexibility of the accelerator, with its wide choice of particle types and energy, and ease of energy changes, combined with an unrivalled collection of spectrometers and polarimeters, made it a splendid tool for unravelling problems in diverse branches of physics. Nevertheless the facility was closed. Let us agree simply to say that the authorities had different priorities to us.

Each of the excellent speakers has had to make a selection from the wealth of material in the subject area of his talk and, in my half hour, I will try to pick out some of the personal highlights of the whole programme of the Laboratory and show how, in many cases, these have influenced developments at other facilities. Though Saturne may be dead, its genes live on!

Before starting this though, let me make one more political point. It is much easier to get finance for doing new experiments at other facilities, than completing the analysis of data taken at Saturne. For the physicists involved it is also more appealing to throw themselves wholeheartedly into the new proposals. However we all know of many experiments, where there are still data on tape representing interesting phenomena, but which risk being left just gathering dust in cupboards. I will not embarrass people by giving examples but I appeal to physicists and their laboratory directors to spend some time and money to extract the maximum from the data taken at Saturne.

⁰Concluding talk at “les 20 ans de Saturne-2” colloquium, Paris, 5 May 1998

It is exactly 30 years since I gave my first talk at Saclay, and in May '68 people had other things on their minds than the application of the Glauber model to the scattering of 1 GeV protons from carbon and oxygen, which was the subject of the talk. Though the Saturne accelerator was completely rebuilt 20 years ago, there has been a continuity in its physics programme and in some of the equipment so that I will stray across the 1977 frontier. Not being a “machine-man”, there is of course a danger that I would not stress sufficiently the role of equipment in my valediction. To avoid this danger, I commend to you the excellent “vulgarisation” of the work carried out at Saturne [1]. Copies of the abridged English translation were given to all participants, although they will be disappointed not to find a description of the bonnes et mauvaises années there!

One should also not forget the Proceedings of the Journées d'Études Saturne, edited over many years by Pierre Radvanyi and Mme. Bordry. These volumes are mines of useful information about Saturne, its equipment and its physics and this is a true reflection of the meetings themselves, which were very important in developing the unique culture of Saturne. As an example, my first ideas on the possibility of quasi-bound $\eta^3\text{He}$ states came to me at the Mont Ste. Odile meeting, and the first calculations were carried out on the train back from Strasbourg to Paris. Other people might remember rather the banquets and the difference between Roscoff and Cavalaire!

Elastic scattering proton-nucleus scattering and the excitation of nuclear levels was the first place where SPESI made an impact on the international stage. As discussed here by Vorobyov, this programme was an extension of the earlier Cosmotron work of Palevsky and friends. Vorobyov's talk reminded me of the calculations of my youth, but I had truly forgotten that the experimental programme of the Gatchina-Saclay collaboration was so vast. Nevertheless it was a programme of classical nuclear physics where the only degrees of freedom were those of the nucleon. Within the Glauber model, they obtained matter densities in many nuclei, and also transition densities, without inserting specific forms for the nuclear densities. Thus they were able to look at changes in density from one isobar to another.

Since Pierre Radvanyi and others have mentioned the influence of Palevsky on the development of Saturne-1, allow me to add a couple of remarks to the history since I worked with Harry in Brookhaven in 1966-67. Saturne-1 was a clone of the Cosmotron, and Harry never understood why the French insisted on taking the original plans for the machine rather than introducing the improvements suggested by the American accelerator theorists who designed the Cosmotron. Secondly, even though Palevsky is now recognised for his work on elastic scattering, his real passion was for the comparison with the inclusive spectra where he hoped to derive nucleon-nucleon correlations from sum rules.

The first really innovative material was an investigation of intermediate energy pick-up reactions, where SPESI could easily measure the angular distributions of $^{12}\text{C}(p, d)^{11}\text{C}^*$ to half a dozen excited states of ^{11}C at 700 MeV [2]. It is fitting that Jacques Thirion presented this at the same conference as Hoistad showed data on the (p, π^+) reaction, since the physics is closely related. Unlike elastic proton scattering

at small angles, these large momentum transfer reactions are intimately related to mesonic or Δ degrees of freedom in the nucleus. This importance of virtual mesons or nuclear isobars, even for reactions where no mesons exist in either the initial or final states, is something that we had to bear in mind in many later experiments at Saturne-2. As a simple rule of thumb, rare events in hadronic reactions tend not to be as rare as would be indicated on the basis of purely nucleonic degrees of freedom.

Personally my most exciting moment at LNS was the night that we measured η production in the $\vec{d}p \rightarrow {}^3\text{He} \eta$ reaction near threshold at SPESIV [3], though it had been studied previously at higher energies [4]. This was also an important moment for the laboratory as well since in this, and subsequent experiments at SPESII [5]:

- It showed a surprisingly high cross section, which is probably due to a virtual pion beam being created on one of the nucleons in the deuteron, with the pion producing an η -meson on the second nucleon. Such ideas are now used in the interpretation of threshold meson production in ion-ion collisions.
- The variation of cross section near threshold implies the existence of a quasibound $\eta {}^3\text{He}$ state, a new form of nuclear matter at high excitation energy, whose existence was clearly confirmed for the $\eta {}^4\text{He}$ system [6, 7].
- The latter provided a natural explanation for the charge symmetry violation signal in the $dd \rightarrow {}^4\text{He} \pi^0$ signal reported by the unforgettable ER54 group just below the η threshold [8].
- It allowed the most precise measurement of the mass of the η meson [9].
- It opened the possibility of studying the rare decays of the η with a tagged beam of such mesons [10]. Important results were found on the decays $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \mu^+\mu^-$ and it is a matter of profound regret that the SPESII programme of rare decay studies had to be cut short for practical reasons. What a catastrophe.
- It lead to the study of heavier continuum states in $pd \rightarrow {}^3\text{He} X$ under threshold conditions [11], as well as the isolation of heavier mesons X such as the ω [12], η' and ϕ [13].
- It also drove people to investigate threshold production of η [14, 15], ω [16], η' [17] and ϕ [18] in nucleon-nucleon scattering using a variety of experimental techniques.

Almost all of these themes, born out of one passionate night of endeavour, will be carried on at other experimental facilities, although the diehard might well say with some justification that we could have done it still better at Saturne! Thus there is a big rare decay programme at Uppsala and searches for quasibound η -nuclear states at Jülich (and during the meeting Paul Kienle explained to me that there were similar proposals at GSI). Both laboratories will carry out production studies of the $\eta {}^3\text{He}$ system and search for charge symmetry violation in $dd \rightarrow {}^4\text{He} \pi^0$. I have

even heard it said that by 2003 François Plouin will have finished his analysis of the mass of the η meson, but I have no confirmation of this.

Saturne was internationally recognised for the intensity and quality of its beams of polarised protons and deuterons [1], which was due to the combination of the Hyperion polarised ion source, the mini-synchrotron Mimas serving as injector, and a meticulous study of depolarising resonances in Saturne itself by the “machinistes”. I was initially confused to find that the “Group Théorie” at Saturne was composed purely of people who studied how the machine worked, whereas there were almost no nuclear/particle theorists in the laboratory. This showed the priorities of the founders of the laboratory and all other machines with which I am familiar have far less backing in this area. Though the successes in Indiana have been outstanding, it has been found that polarised particles are difficult to accelerate at both Uppsala and Jülich. In his talk Lagniel highlighted the collaborations between machinistes and experimentalists, especially those of nucleon-nucleon, and showed how both sides benefitted from this. It wasn’t always war!

The major user of polarised protons was the nucleon-nucleon group and often in the Comité des Expériences we exploited the fact that there was always a colleague who was willing to take beam time during any holiday period. It might be a coincidence but the moment that they finished their programme on proton-proton and proton-neutron elastic scattering, the Authorities closed the Laboratory down. The nucleon-nucleon studies represented perhaps the most fundamental of the research that was carried out at Saturne. This was done in a completely professional manner and in many cases the group had sufficient data in order to make direct reconstructions of the amplitudes without passing through theoretical models.

Where Saturne was really unique was in respect of its deuteron beams, which combined high intensity with large values of both vector and tensor polarisation. Without Saturne we could not have calibrated the AHEAD [19] or POLDER [20] deuteron tensor polarimeters which, as Garçon described in his talk, played vital roles in separating the deuteron form factors through measurements of the polarisation of the recoil deuteron in elastic electron-deuteron scattering. The Comité des Expériences and Management of Saturne always took the broad view that the advancement of Physics was the central theme of the Laboratory and, if this involved calibrating equipment for good experiments to be done elsewhere, then this was a price worth paying. It certainly paid off in the case of the deuteron form factor. Michel Garçon has shown us the first preliminary results from CEBAF on the value of T_{20} in elastic $ed \rightarrow e'd'$. The collaboration could work to values of q^2 twice as big as earlier experiments with a much bigger efficiency than all competing techniques.

The POLDER polarimeter is based upon the deuteron charge exchange reaction $\vec{d}p \rightarrow \{pp\}n$, which was predicted to show a strong analysing power signal if the final proton-proton pair had small excitation energy and was produced with a small momentum transfer relative to the incident deuteron [21]. This reaction was investigated in the few hundred MeV range [22] but also in the GeV range through the remarkable technique of detecting both protons in the focal plane of the SPESIV spectrometer [23]. This was one of a whole series of charge-exchange experiments

with different probes undertaken over years by the Copenhagen-Orsay group, and it is a matter of profound regret that the group leader, Carl Gaarde, was taken from us a few weeks before this meeting. Michelle Roy-Stéphan has reminded us of the enormous quantity of results, especially in the ($^3\text{He}, t$) reaction, achieved over 14 years and the successes in their interpretation has come through active collaborations with theorists. One of these, Madeleine Soyeur, showed how such results could be exciting for theorists and I believe that she communicated well this enthusiasm to the machinistes in the hall, even if they couldn't appreciate some of the finer points.

The method of detecting two protons in SPESIV could be extended to looking also at deep inelastic scattering, where a Δ -isobar had been produced on hydrogen or nuclear targets [24]. The $(\vec{d}, 2p)$ reaction is an interesting probe for $\Delta T = \Delta S = 1$ transitions but, inspired by Marcel Morlet, the Orsay group showed that measurements of vector spin transfer in $(\vec{d}, \vec{d}')$ can provide useful signals for $\Delta T = 0, \Delta S = 1$ states in the residual nucleus [25]. This needed a lot of intuition because the symmetry used here is only an approximate one and a complete separation would have required the combination of a tensor polarised beam and tensor polarimeter.

Of equally lasting importance was the whole series of measurements designed to investigate the few body problem in intermediate energy nuclear physics. The measurement of triple spin parameters in $\vec{d}\vec{p} \rightarrow \vec{p}d$ at 1.6 GeV [26] represented a real tour de force which is never likely to be repeated. More appealing are however the survey experiments which measure a single observable, such as the deuteron tensor analysing power T_{20} , as a function of beam energy, as was done for $\vec{d}p \rightarrow pd$ in the backward direction [27] or pion production through $\vec{d}p \rightarrow ^3\text{He}\pi^0$ [28, 29]. Due to its D-state component, the deuteron has a geometric deformation looking in momentum space a bit like a pancake. This could be investigated by measuring the T_{20} dependence of the Fermi momentum in deuteron break up $\vec{d}p \rightarrow ppn$ [30] but the wealth of polarimeters at Saturne also permitted a polarisation measurement of a final proton [31].

The D-states in ^6Li could be studied in analogous break-up measurements [32] with beams of polarised ions of several GeV/c (which must be high compared to the Fermi momentum), and this seems to have been the only published experiment using such a beam. The MIMAS synchrotron was designed to furnish high quality polarised beams and also beams of ions up to ^{129}Xe (30^+) but the demand for such heavy ions has represented only about 10% of the total requests over many years, and even less of the publications. In part this might be that the competition from other facilities in the heavy ion field, such as Ganil or SIS, has proved more intense than for the light ions where Saturne was supreme. However, apart from Diogène, there were no specific heavy ion detectors at Saturne and groups tended to bring small equipment with them for multifragmentation or other studies [33]. Saturne was the great success that it was for light ions because it had a whole collection of outstanding spectrometers which were tailored for the purpose.

I have already stressed that physics must be exciting and no survey of the work at Saturne would be complete without mentioning “Le pari de Pascal” and the human desire to find something really revolutionary, even if the probability of success were

infinitesimally small. If there were a good bridge player in the room, then he would tell you that you must think defensively, such that the international competitors should never be allowed to make too good a score. Thus Saturne could not allow other laboratories to make the earth-shattering discoveries which might have been made at Saclay. In other fields this *maladie* took the form of trying to repeat the discoveries of Pons and Fleischmann on cold fusion, but at Saturne it was the hunt for narrow dibaryon resonances [34], pion-nucleon bound states [35], abnormal production of pions [36], the anomalous [37] *etc.* It was really like something out of “Les vacances de M. Hulot”. Nothing convincing was ever found which was not at the limits of systematic/statistical error bars. Colleagues will be relieved to know that this approach is being carried on in the successor laboratories and that I was equally unsuccessful in stopping the search for the d' at COSY and CELSIUS PAC's as I was with the narrow dibaryons at the Saturne Comité des Expériences.

It is true that just one such discovery might have provided some defence for Saturne and it is often these searches which bring out the best in technical ingenuity. One example of this is the famous wheel of Beurtey and Saudinos [38]. In order to make simultaneous measurements of A_y in proton-proton elastic scattering at many different energies, they constructed an energy degrader with many steps, looking something like a circular Escher staircase. By letting this rotate fast in the beam, high statistics data could be obtained at 16 different energies simultaneously. A short run was sufficient to kill off one dibaryon (but not French).

Over the years Saturne has been invaluable for the calibration of much other equipment in addition to polarimeters. It is true that Saturne will not be much remembered for such mundane work in the publications of physics carried out some time later at other laboratories. However people will remember Saturne when they find that they can no longer use it to test counters for LHC for example. Often the applied research at Saturne gave rise to interesting physics in related fields, and I have in mind here the some of the work described by Rolf Michel on pseudo-meteorites.

You will have noticed that I have not spoken at all about the three big programmes which have dominated the last one or two years of the running of Saturne-2, *viz* DISTO, SPESIV π , and Nuclear Transmutation; it is much too early to judge their significance. We all of course hope that the information provided by the transmutation experiments will prove valuable in a search for a sensible method for dealing with nuclear waste. That really would be a worthwhile monument for Saturne to compete with the statue of the Ptolemy which was unveiled today outside the Grand Palais.

Pour terminer, je dois remercier Alain, Françoise, Simone et Bernard qui ont assuré le succès de cette réunion. Il est bien évident que le français n'est pas la langue maternelle, ni de M. Blair ni de moi. Néanmoins il va rester ma langue fraternelle et cela est d'une grande part dû aux collaborations fructueuses avec les gens autour de Saturne, beaucoup trop nombreux à mentionner.

Saturne, nous allons tous nous souvenir de vous. Saturne, adieu! Merci!

References

- [1] R. Beurtey and A.-M. Gauriot, *De la Technique à la Physique des Energies Intermédiaires*, (Laboratoire National Saturne, 1987).
- [2] J. Thirion, in *High Energy Physics and Nuclear Structure*, ed. G. Tibell (North Holland, 1973).
- [3] J. Berger *et al.*, *Phys. Rev. Lett.* **61**, 919 (1988).
- [4] P. Berthet *et al.*, *Nucl. Phys. A* **443**, 589 (1985).
- [5] B. Mayer *et al.*, *Phys. Rev. C* **53**, 2068 (1996).
- [6] R. Frascaria *et al.*, *Phys. Rev. C* **50**, R537 (1994).
- [7] N. Willis *et al.*, *Phys. Lett. B* **406**, 14 (1997).
- [8] L. Goldzahl *et al.*, *Nucl. Phys. A* **533**, 675 (1991).
- [9] F. Plouin *et al.*, *Phys. Lett. B* **276**, 526 (1992).
- [10] R. Abegg *et al.*, *Phys. Rev. D* **50**, 92 (1994).
- [11] F. Plouin in *Production and Decay of Light Mesons*. ed. P. Fleury (World Scientific, Singapore, 1989).
- [12] R. Wurzinger *et al.*, *Phys. Rev. C* **51**, R443 (1993).
- [13] R. Wurzinger *et al.*, *Phys. Lett. B* **374**, 283 (1996).
- [14] A.M. Bergdolt *et al.*, *Phys. Rev. D* **48**, R2969 (1993).
- [15] E. Chiavassa *et al.*, *Phys. Lett. B* **337**, 192 (1994)).
- [16] F. Hibou *et al.*, (in preparation).
- [17] F. Hibou *et al.*, *Phys. Lett. B* (in press).
- [18] R. Bertini, spokesman for the DISTO collaboration (private communication), and W. Kühn (this meeting).
- [19] J.M. Cameron *et al.*, *Nucl. Instrum. Methods A* **305**, 257 (1991).
- [20] S. Kox *et al.*, *Nucl. Instrum. Methods A* **346**, 527 (1994).
- [21] D.V. Bugg and C. Wilkin, *Phys. Lett. B* **152**, 37 (1985).
- [22] S. Kox *et al.*, *Nucl. Phys. A* **556**, 621 (1993).
- [23] C. Ellegaard *et al.*, *Phys. Rev. Lett.* **59**, 974 (1987).

- [24] T. Sams *et al.*, *Phys. Rev. C* **51**, 1945 (1995).
- [25] B.N. Johnson *et al.*, *Phys. Rev. C* **51**, 1726 (1995).
- [26] V. Ghazikhanian *et al.*, *Phys. Rev. C* **43**, 1532 (1991).
- [27] J. Arvieux *et al.*, *Nucl. Phys. A* **431**, 613 (1984).
- [28] C. Kerboul *et al.*, *Phys. Lett. B* **181**, 28 (1986).
- [29] V.N. Nikulin *et al.*, *Phys. Rev. C* **54**, 1732 (1996).
- [30] V. Punjabi *et al.*, *Phys. Rev. C* **39**, 608 (1989).
- [31] S.L. Belostotski *et al.*, *Phys. Rev. C* **56**, 50 (1997).
- [32] V. Punjabi *et al.*, *Phys. Rev. C* **46**, 984 (1992).
- [33] C. Williams *et al.*, *Phys. Rev. C* **55**, R2132 (1997).
- [34] B. Tatischeff *et al.*, *Phys. Rev. C* **36**, 1995 (1987).
- [35] B. Tatischeff *et al.*, *Phys. Rev. Lett.* **79**, 601 (1997).
- [36] J. Julien *et al.*, *Z. Phys. A* **347**, 181 (1994).
- [37] M. Bedjidian *et al.*, *Z. Phys. A* **327**, 337 (1987).
- [38] R. Beurtey *et al.*, *Phys. Lett. B* **293**, 27 (1992).